

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



P 12515 ET

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 7/00, 7/40, 7/42, 7/44, 31/445</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/41826</b>
			<b>(43) International Publication Date:</b> 13 November 1997 (13.11.97)
<b>(21) International Application Number:</b> PCT/US97/07699			<b>(81) Designated States:</b> AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
<b>(22) International Filing Date:</b> 6 May 1997 (06.05.97)			
<b>(30) Priority Data:</b> 60/016,769 7 May 1996 (07.05.96) US			
<b>(71) Applicant (for all designated States except US):</b> THOMAS JEFFERSON UNIVERSITY [US/US]; 11th and Walnut Streets, Philadelphia, PA 19107 (US).			
<b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> BERNSTEIN, Eric [US/US]; 1321 Grenox Road, Wynnewood, PA 19096 (US).			
<b>(74) Agents:</b> LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).			<b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

**(54) Title:** USE OF TEMPOL IN THE PREVENTION OF PHOTOAGING

**(57) Abstract**

A method of preventing photoaging and other types of sun damage by topically applying a composition containing Tempol is provided. Pharmaceutical compositions comprising Tempol for the prevention of photoaging and other types of sun damage are also provided.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MX	Mexico	UA	Ukraine
BR	Brazil	IL	Israel	MY	Malaysia	UG	Uganda
BY	Belarus	IS	Iceland	NE	Niger	US	United States of America
CA	Canada	IT	Italy	NL	Netherlands	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NO	Norway	VN	Viet Nam
CG	Congo	KE	Kenya	NZ	New Zealand	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	PL	Poland	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PT	Portugal		
CM	Cameroon	KR	Republic of Korea	RO	Romania		
CN	China	KZ	Kazakhstan	RU	Russian Federation		
CU	Cuba	LC	Saint Lucia	SD	Sudan		
CZ	Czech Republic	LI	Liechtenstein	SE	Sweden		
DE	Germany	LK	Sri Lanka	SG	Singapore		
DK	Denmark	LR	Liberia				
EE	Estonia						

## USE OF TEMPOL IN THE PREVENTION OF PHOTOAGING

## BACKGROUND OF THE INVENTION

The effects of ultraviolet radiation from exposure to the sun on human skin are a growing concern for today's longer-lived population. The majority of changes associated with an aged appearance result from chronic sun-damage. Warren et al., *J. Am. Acad. Dermatol.* 1991, 25:751-760; Frances, C. and Robert, L., *Int. J. Dermatol.* 1984, 23:166-179. Dramatic alterations of the superficial dermis accompany the deep wrinkles and laxity common in photoaged skin. The major histopathologic alteration of photoaged skin is the accumulation of material which, on routine histopathologic examination, has the staining characteristics of elastin and is, thus, termed solar elastosis. Immunohistochemical staining has shown the poorly-formed fibers comprising solar elastosis to be composed of elastin (Chen et al., *J. Invest. Dermatol.* 1986, 87:334-337; Mera et al., *Br. J. Dermatol.* 1987, 117:21-27) fibrillin (Chen et al., *J. Invest. Dermatol.* 1986, 87:334-337; Dahlback et al., *J. Invest. Dermatol.* 1990, 94:284-291; Bernstein et al., *J. Invest. Dermatol.* 1994, 103:182-186) and versican, the normal components of elastic fibers (Zimmerman et al., *J. Cell. Biol.* 1994, 124:817-825). A coordinate increase in elastin, fibrillin and versican mRNAs has been demonstrated in fibroblasts derived from photodamaged skin, as compared to fibroblasts derived from normal skin from the same individuals. Bernstein et al., *J. Invest. Dermatol.* 1994, 103:182-186. Elevated elastin mRNA levels in sun-damaged skin result from enhanced elastin promoter activity, as shown by transient

transfections of fibroblasts with a DNA construct composed of the human elastin promoter linked to the chloramphenicol acetyltransferase (CAT) reporter gene. Bernstein et al., *J. Invest. Dermatol.* 1994, 103:182-186.

5 The generation of free radicals following exposure of the skin to ultraviolet radiation is well known in the art. Free radical mechanisms have been shown to be responsible for redness and erythema resulting from exposure to ultraviolet radiation. A number of antioxidants have been tested as  
10 photoprotective agents, however, results from these studies indicate that the ability of these agents to provide protection is variable.

Miyachi Y., *J. Dermatol. Sci.* 1995, 9:79-86 provides a review of photoaging from a photo-oxidative standpoint and  
15 suggests the use of antioxidants as regulators of photoaging. Specifically, studies with superoxide dismutase (SOD) are described. However, it is concluded that sunscreen agents provided better protection from ultraviolet radiation.

Bissett et al. *Photodermatol. Photoimmunol. Photomed.*  
20 1990, 7:56-62 demonstrated that mice, topically treated with solutions of superoxide-scavenging anti-oxidants such as alpha-tocopherol, ascorbic acid, propyl galate and Trolox prior to ultraviolet B (UVB) radiation exposure, exhibited significantly less damage than untreated mice. However, additional  
25 antioxidants or free radical scavengers that were tested, including glutathione, beta-carotene, BHT, mannitol, divinylglycol, pantetheine, urea and histidine, provided no significant protection against UVB radiation. Further, the severity of UVA radiation-induced mouse skin damage was not  
30 reduced by topical application of these antioxidants in these studies. Thus, it is clear that the current approaches used to prevent the cumulative effects of photoaging are inadequate.

Historically, more research has been done in the area of radiation oncology. Damage from ionizing radiation and a  
35 portion of ultraviolet radiation-induced damage has been shown to be due to the formation of radical oxygen species. Sulfhydryl compounds were among the first radioprotectors to be

identified. Their protective mechanism appears to be due to their ability to scavenge radiation-induced free radicals and/or donate reducing equivalents to oxidized molecules. Hematopoietic cytokines have also been investigated as radioprotectors. They are believed to protect by more quickly restoring hematopoietic function after radiation exposure.

Recently, a new class of radioprotectors, the nitroxides, has been described. As a class, nitroxides are stable free radical components which react with a variety of biologically relevant compounds including other free radicals (Nilsson et al. *J. Biol. Chem.* 1989, 264:11131-11135). The observation that several nitroxides themselves reacted with free radicals, specifically oxy radicals, led to the investigation of these compounds as radioprotectors (Samuni et al. *Free Radical Biol. Med.* 1989, 6:141-148).

Tempol [4-hydroxy-2,2,6,6-tetramethyl-piperidinyloxy, free radical] is a piperidinyloxy with the n-oxide sterically stabilized by symmetric pairs of adjacent methyl groups. This compound is commercially available through Aldrich Chemical Co., Milwaukee, WI. It is most commonly used to spin label biological molecules such as NADP.

Tempol has been demonstrated to function as a superoxide dismutase (SOD) mimic, protecting mammalian cells from superoxide generated from hypoxanthine/xanthine oxidase and from hydrogen peroxide mediated cytotoxicity (Mitchell et al. *Biochem.* 1990, 29:2802-2807; Samuni et al. *J. Biol. Chem.* 1988, 263:17921-17924). Tempol has also been demonstrated to provide both *in vitro* and *in vivo* protection against ionizing radiation (Mitchell et al. *Arch. Biochem. Biophys.* 1991, 289:62-70). Recently, Tempol was shown to protect against radiation induced alopecia by speeding the recovery of hair growth within a field of heavily irradiated skin (Goffman et al. *Int. J. Rad. Onc. Biol. Phys.* 1992, 22:803-806). Goffman et al. suggest that this protection is due to direct protection of hair follicle stem cells and development of other nitroxides. Importantly, this study also demonstrated that Tempol is undetectable in the blood following application. This is of concern because

- 4 -

topical radioprotectors share a tendency towards systemic absorption, raising the potential problem of tumor radioprotection.

It has now been found that topical application of Tempol prevents photoaging and other skin damage resulting from exposure to ultraviolet radiation.

#### SUMMARY OF THE INVENTION

In the present invention, a new use is provided for the compound Tempol (4-hydroxy-TEMPO, free radical; 4-hydroxy-2,2,6,6-tetramethylpiperidinyloxy, free radical). It has now been found that Tempol prevents photoaging and blocks UVB thereby preventing other sun-damage such as sunburn or skin cancer. In addition, Tempol has now been shown to protect against UVA in vitro. This compound not only functions as a free-radical scavenger, but also possesses sun blocking ability. The ability to block ultraviolet radiation-induced damage has been demonstrated. Compositions for use as sunscreen agents comprising Tempol are also provided.

#### DETAILED DESCRIPTION OF THE INVENTION

Profound changes take place in the superficial dermis as a result of chronic sun-exposure. The major alteration is the deposition of massive amounts of abnormal elastic material, termed solar elastosis. It has been shown that solar elastosis is accompanied by elevations in elastin and fibrillin mRNAs and elastin promoter activity.

A transgenic mouse model which contains the human elastin promoter linked to a chloramphenicol acetyltransferase (CAT) reporter gene for testing compounds that may inhibit cutaneous photodamage has been developed. These mice express human elastin promoter activity in a tissue-specific and developmentally regulated manner. Promoter activity can be studied in this model as a function of small increases in ultraviolet radiation, demonstrating the sensitivity of the assay. In addition, quantitative data can be obtained after only a single exposure to ultraviolet radiation. A test

- 5 -

compound is applied to the skin of a transgenic mouse capable of expressing the human elastin promoter. The transgenic mouse is then exposed to ultraviolet radiation and human elastin promoter activity in the mouse is determined. The human  
5 elastin promoter activity is then compared to that in transgenic mice also exposed to an equivalent dose of ultraviolet radiation which were not treated with the test compound to determine whether or not the test compound provided protection against the ultraviolet radiation. Since elastin  
10 promoter activation is a primary event in cutaneous aging, these mice represent a mouse model of human photoaging.

Using this transgenic mouse line, the ability of Tempol to inhibit the effects of ultraviolet radiation on human elastin promoter activity was determined. In these  
15 experiments, 4-5 day old mice received either no treatment, 0.4M Tempol, 245 mJ/cm<sup>2</sup> UVB, or 245 mJ/cm<sup>2</sup> UVB and 0.4 M Tempol. Tempol was diluted in ethanol and was applied topically to the backs of these mice. Following phototreatment, the backs of the mice were rinsed twice with 70% isopropyl alcohol pads to  
20 remove any excess Tempol. Mice were sacrificed and skin harvested for determination of CAT activity 24 hours after phototreatment. Tempol alone produced no significant change, raising CAT activity 1.7 fold (s.d. = 0.48). UVB increased CAT activity 7.2-fold (standard deviation (s.d.) = 0.72), and the  
25 addition of Tempol to UVB-treated mice decreased the elevation to 3.3-fold (s.d.= 0.68) that of untreated controls.

Experiments demonstrating the ability of Tempol to protect against UVB-induced promoter activation were also performed in fibroblast cultures established from the skin of  
30 transgenic mice. Cells received either no Tempol or UVB (controls), 50 mM Tempol alone without UVB, 5.5 mJ/cm<sup>2</sup> UVB alone, pretreatment with 25 mM Tempol and subsequent exposure to 5.5 mJ/cm<sup>2</sup> UVB, or pretreatment with 50 mM Tempol and subsequent exposure to 5.5. mJ/cm<sup>2</sup> UVB. Tempol was added to  
35 culture dishes ten minutes prior to exposure to UVB and left on the cells 4 minutes after UVB exposure. Tempol alone did not alter CAT activity in these cells to a significant degree. UVB

- 6 -

alone increased CAT activity to 9.5-fold (s.d. = 0.47) that of unirradiated cells. The addition of 25 mM Tempol to UVB-treated cells decreased the elevation to 4.2-fold (s.d. = 0.02) that of controls, while 50 mM Tempol further decreased the  
5 elevation to 2.8-fold (s.d. = 1.4) that of controls.

Additional experiments were performed in cells with varying amounts of UVB. In these experiments, control cells received no UVB or Tempol. UVB controls received either 0.7, 1.4, 2.7 or 5.5 mJ/cm<sup>2</sup> of UVB. Tempol controls received 100 mM  
10 of Tempol and no UVB. UVB treated cells were preincubated with phosphate buffered saline (PBS) or 100 mM Tempol 12.5 minutes before UVB. The effect of UVB and Tempol on cell viability was determined using a trypan blue exclusion kit.

Before application to cultures, tissue culture medium was  
15 removed and the cells were rinsed twice with PBS. Cells were incubated in 10 cm diameter dishes with 3 ml of 100 mM Tempol or PBS for 12.5 minutes prior to UVB exposure. UVB controls were placed in an equal amount of PBS 12.5 minutes prior to irradiation with UVB. After treatment, cells were cleansed  
20 twice with PBS and the tissue culture medium was replaced in all dishes. Cells were harvested for determination of CAT activity 24 hours after phototreatment. Only fibroblasts from mice representing the same litter were used for any given experiment, and utilized in passage two to three times. Two  
25 dishes of cells were used for each data point, and experiments were repeated in triplicate. Thus, six values were determined for each experimental condition. For statistical analysis, a paired t-test was performed. Data from these experiments is depicted in Table 1.



TABLE 1: In Vitro Effect of Tempol on UVB-induced induction of CAT activity

UVB Dose (mJ/cm <sup>2</sup> )	UVB CAT	UVB + Tempol CAT	% Protection from UVB
0.7	3.2	1.9	41 p<0.01
1.4	8.1	2.1	74 p<0.001
2.8	15.7	4.8	69 p<0.001
5.5	20.0	11.6	42 p<0.001

As in previous experiments, Tempol alone did not alter CAT activity to a significant degree (1.2-fold that of untreated cells). UVB alone increased CAT activity 3.2-, 8.1-, 15.7- and 20.0-fold that of control cells for UVB doses of 0.7, 1.4, 2.7 and 5.5 mJ/cm<sup>2</sup> of UVB, respectively. The addition of 100 mM Tempol to UVB-treated cells decreased this elevation to 1.9-, 2.1-, 4.8- and 11.6-fold greater than untreated controls for UVB doses of 0.7, 1.4, 2.7 and 5.5 mJ/cm<sup>2</sup> of UVB, respectively.

The free radical scavenging ability of Tempol was also demonstrated in cells. Cells received either no treatment, 5.5 mJ/cm<sup>2</sup> of UVB, UVB with 50 mM Tempol suspended in a quartz dish above the cells covered in saline, or UVB with 50 mM Tempol applied directly to the cells with saline in a quartz dish suspended above the cells. Tempol suspended above the cells functions as a UVB blocker. Tempol applied directly to the cells functions as both a UVB blocker and a scavenger of free radicals. UVB alone resulted in a 14.2-fold (s.d. = 1.5) increase in CAT activity. Tempol suspended above the cells in a quartz dish (and thus acting as a screen only) resulted in a 2.4-fold increase in CAT activity as compared to untreated controls, which is almost a 6-fold reduction from cells treated with UVB alone and not receiving Tempol, thus demonstrating significant protection from UVB. When Tempol was placed in direct contact with cells, CAT activity was reduced to 1.5-fold that of controls. Accordingly, the scavenging properties of Tempol reduced CAT activity a further 38%. Thus, in addition to providing protection against photoaging by absorbing light,

- 8 -

Tempol has also been demonstrated to offer protection against damage once ultraviolet radiation has already reached the skin.

Experiments demonstrating the ability of Tempol to protect against UVA-induced promoter activation were also performed in fibroblast cultures established from the skin of transgenic mice. Because a response to UVA alone *in vitro* is not measurable and the *in vivo* response is relatively small, 8-methoxypsoralen (8-MOP) was added to the system to enhance the promoter response to UVA. The addition of 8-MOP resulted in a substantial increase in CAT activity following administration of UVA, making this model more useful for studying agents which may protect against UVA as well as UVB.

In these experiments, control cells received no UVA or 8-MOP. UVA controls received either 0.5, 0.75 or 1.0 J/cm<sup>2</sup> of UVA without prior incubation with 8-MOP. 8-MOP controls received 1 µg/ml of 8-MOP but no UVA. Tempol controls received 100 mM Tempol and no UVA or 8-MOP. PUVA-treated cells were preincubated with 1.0 µg/ml of 8-MOP in PBS and then exposed to 0.5, 0.75 or 1.0 J/cm<sup>2</sup>. Tempol treated cells were pretreated with 100 mM Tempol 12.5 minutes prior to UVA exposure. 8-MOP was also added to these cells at the same time as the Tempol.

It was found that the combination of 1 µg/ml of 8-MOP and 0.5, 0.75 or 1.0 J/cm<sup>2</sup> of UVA in fibroblast cultures resulted in increases in CAT activity to 2.7-, 9.0- and 21.4-fold that of controls, respectively. UVA alone and 8-MOP alone failed to produce significant CAT activity. The addition of Tempol to the cultures reduced the PUVA-induced increase in CAT activity to 1.5-, 1.2- and 2.0-fold that of controls, respectively, thus almost completely eliminating the PUVA-induced increase in CAT activity. The absorption spectrum of Tempol has shown very little absorption in the UVA range, making the protection by Tempol against PUVA-induced elastin gene induction almost entirely due to the free radical scavenging abilities of Tempol. In addition, suspending Tempol above the cells failed to afford protection against the PUVA-induced increases in CAT activity, thus further demonstrating that protection by Tempol against UVA-induced photodamage is almost entirely due to its

- 9 -

free radical scavenging properties.

Accordingly, topical application of a composition comprising Tempol to the skin provides protection against photoaging and other sun-damage such as sunburn and skin cancer. Examples of compositions comprising Tempol include, but are not limited to creams, lotions and sprays. Methods of formulating Tempol into creams, lotions and sprays as well as pharmaceutical additives for such formulations are well known to those skilled in the art. As will be obvious to those skilled in the art upon this disclosure, such compositions may further comprise secondary or additional sunscreens or free radical scavengers such as, but not limited to, Vitamin C and Vitamin E and analogs thereof. In a preferred embodiment, a composition comprising Tempol is applied to the skin prior to exposure to the sun. However, application of these compositions subsequent to the exposure can also mitigate any damage resulting to the skin from this exposure. It is believed that these compositions of the present invention will be especially useful in protecting individuals with heightened sensitivities to the sun, such as, but not limited to, individuals undergoing psoralen treatment for cancer, psoriasis and other skin conditions; individuals undergoing photodynamic therapy for skin cancer, psoriasis and other skin conditions; individuals suffering from genetic repair defects such as xeroderma pigmentosa, albinism or other conditions resulting from decreased endogenous melanin pigment.

The following nonlimiting examples are provided to further illustrate the present invention.

- 10 -

**EXAMPLES****Example 1: Transgenic mice expressing the human elastin promoter**

A homozygous line of transgenic mice expressing the 5.2-  
5 kb human elastin promoter linked to a CAT reporter gene was  
used. Hsu-Wong et al., *J. Biol. Chem.* 1994, 269:18072-18075.  
These mice express the human elastin promoter in a tissue-  
specific and developmentally regulated manner. Mice four or  
five days old were used since at this age, visible hair growth  
10 is not yet present.

**Example 2: Fibroblast Cultures**

Fibroblast cultures were established from the skin of  
transgenic mice by explanting tissue specimens onto plastic  
tissue culture dishes and allowing cells to migrate to the  
15 surrounding area. The primary cultures were maintained in  
Dulbecco's modified Eagle's medium supplemented with 10% fetal  
calf serum, 2 mM L-glutamine and antibiotics, at 37°C. The  
primary cell cultures were passaged by trypsinization, and the  
subcultures in passages three or four were utilized for UVB  
20 treatment. After UV exposure, the cells were incubated in  
tissue culture medium for 24 hours, and then harvested for  
determination of CAT activity.

**Example 3: CAT Assay**

25 To measure the expression of the human elastin  
promoter/CAT reporter gene construct in the skin of transgenic  
mice and in fibroblast cultures established from these animals,  
CAT activity was determined. For extraction of the CAT from  
skin, the specimens were homogenized in 0.25 Tris-HCl, pH 7.5,  
30 using a tissue homogenizer (Brinkmann Instruments, Inc.  
Westbury, NY). The homogenates were centrifuged at 10,000 X g  
for 15 minutes at 4°C and the protein concentration in the  
supernatant determined by a commercial protein assay kit (Bio-  
Rad Laboratories, Richmond, CA). Aliquots of the supernatant  
35 containing 100 µg of protein were used for assay of CAT  
activity by incubation with [<sup>14</sup>C] chloramphenicol in accordance

- 11 -

with well-known procedures. The acetylated and non-acetylated forms of radioactive chloramphenicol were separated by thin-layer chromatography and CAT activity was determined by the radioactivity in the acetylated forms as a percent of the total  
5 radioactivity in each sample.

**Example 4: UVB Sources**

Irradiation with UVB for *in vitro* and *in vivo* studies were performed using a closely spaced array of seven Westinghouse FS-40 sunlamps, which delivered uniform  
10 irradiation at a distance of 38 cm. The energy output at 38 cm was measured with a Solar Light Model 3D UVA and UVB detector (Solar Light Company, Philadelphia, PA). The output of FS-40 sunlamps was 23.4 units/hour of UVB at 38 cm, where each unit is equivalent to 21 mJ/cm<sup>2</sup> or erythema effective energy.

**15 Example 5: UVA Sources**

Irradiation with UVA was performed using a closely spaced array of seven Sylvania FR40T12 PUVA lamps filtered through window glass of 2 mm thickness to remove wavelengths below 320 nm. The energy output at 38 cm from the array was measured  
20 with a Solar Light Model 3D UVA and UVB detector (Solar Light Company, Philadelphia, PA). The output of the lamps filtered through the window glass was 2.02 mW/cm<sup>2</sup>, with no detectable UVB.

- 12 -

**What is Claimed:**

1. A method of preventing photoaging and other sun-damage comprising topically applying to the skin a composition containing Tempol.
- 5       2. The method of claim 1 wherein said composition is applied prior to exposure of the skin to the sun.
3. The method of claim 1 wherein said composition is applied subsequent to exposure of the skin to the sun.
- 10       4. A method of protecting individuals with a heightened sensitivity to the sun from damage resulting from the sun comprising topically applying to the skin of an individuals with a heightened sensitivity to the sun a composition containing Tempol prior to exposure of the individual to the sun.
- 15       5. A pharmaceutical composition for prevention of photoaging and other sun-damage comprising Tempol and a pharmaceutical additive.
6. The pharmaceutical composition of claim 5 further comprising a secondary sunscreen or free radical scavenger.

## INTERNATIONAL SEARCH REPORT

 International application No.  
 PCT/US97/07699

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 7/00, 7/40, 7/42, 7/44, 31/445

US CL : 424/59, 60, 400, 401; 514/315

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/59, 60, 400, 401; 514/315

 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
 NONE

 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
 APS, CAS ONLINE: PHOTOAGING, TEMPOL, COSMETIC, PHARMACEUTICAL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	US 5,569,663 A (RIBIER et al.) 29 October 1996, see entire document.	1-6
A	US 5,462,946 A (MITCHELL et al.) 31 October 1995, see entire document.	1-6

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\*

document member of the same patent family

Date of the actual completion of the international search

01 OCTOBER 1997

Date of mailing of the international search report

15 OCT 1997

 Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT

Authorized officer

STEPHEN A. DODSON



